



**NATIONAL COMMISSIONER OF THE ICELANDIC POLICE**  
DEPARTMENT OF CIVIL PROTECTION AND EMERGENCY MANAGEMENT



**THE SCIENTIFIC ADVISORY BOARD OF THE ICELANDIC CIVIL PROTECTION**

**Date: 26.09.2014**

**Time: 09:30**

**Location: Crisis Coordination Centre, Skogarhlid.**

**Regarding: Volcanic activity in the Bardarbunga system.**

**Attending: Scientists from Icelandic Met Office and the Institute of Earth Sciences University of Iceland along with representatives from the Icelandic Civil Protection and the Directorate of Health.**

**Main points**

- **Volcanic eruption in Holuhraun**
- **Air quality**
- **Scenarios**

**Notes**

- The new lava field continues to grow and has now crossed the track (in Flæður) leading into the Holuhraun area.
- The subsidence of the Bardarbunga caldera continues with same rate as before.
- Seismic activity in Bardarbunga continues on similar rate as the last few days. Five earthquakes bigger than M3,0 were recorded since noon yesterday. The biggest one was M5,0 at 16:35 yesterday afternoon.
- Smaller earthquakes were detected in north part of the dyke and around the eruption site.
- GPS measurements show continuing slow movements.
- No change was detected in water monitoring that cannot be explained with changing weather.

**Air quality:**

- Yesterday a high concentration of SO<sub>2</sub> was measured around lake Myvatn (2000 microgram pr. cubic meter) and last night in Reydarfjordur (2600 microgram pr. cubic meter). The Environmental Agency of Iceland is waiting for shipment of SO<sub>2</sub> meters that will be put up around Iceland.
- Pollution from the eruption is mostly expected to move towards east and southeast today (Friday) and tomorrow. A map showing the gas forecast can be found on the web page of the Icelandic Met Office [www.vedur.is/vedur/spar/textaspar/oskufok/](http://www.vedur.is/vedur/spar/textaspar/oskufok/) An interactive map showing the gas distribution can be seen at [www.vedur.is/vedur/spar/gasdreifing](http://www.vedur.is/vedur/spar/gasdreifing)
- The Icelandic Met Office has also opened a web page where people can report if they have detected gas pollution. A link to the page can be found on the Icelandic version of the web page under [Skrá mengun](#).
- **Instructions:**
  - People who feel discomfort are advised to stay indoors, close their windows, turn up the heat and turn off air conditioning. Use periods of good air quality to ventilate the house. Measurements of air quality can be found on the webpage [www.loftgaedi.is](http://www.loftgaedi.is) The Meteorological Office issues forecast on its web-page and warnings if conditions change to the worse.
  - Instructions from the office of the Chief Epidemiologist and The Environment Agency can be found on their web-sites [www.ust.is](http://www.ust.is) and [www.landlaeknir.is](http://www.landlaeknir.is)
  - The Icelandic Met Office will publish forecasts for sulphuric gases dispersion on the web and in the national radio. It will also be endeavored/seeked to broadcast the forecasts on national television.



# NATIONAL COMMISSIONER OF THE ICELANDIC POLICE

## DEPARTMENT OF CIVIL PROTECTION AND EMERGENCY MANAGEMENT



- Information and any questions on air pollution can be sent to The Environment Agency through the email [gos@ust.is](mailto:gos@ust.is). The Environment Agency is especially looking for information from people who have been in contact with high concentrations of gas; where they were, at what time it happened, how the gas cloud looked (colour and thickness of the cloud) and how they were affected by it.
- Three scenarios are considered most likely:
  - The eruption on Holuhraun declines gradually and subsidence of the Bardarbunga caldera stops.
  - Large-scale subsidence of the caldera occurs, prolonging or strengthening the eruption on Holuhraun. In this situation, it is likely that the eruptive fissure would lengthen southwards under Dyngjufokull, resulting in a jokulhlaup and an ash-producing eruption. It is also possible that eruptive fissures could develop in another location under the glacier.
  - Large-scale subsidence of the caldera occurs, causing an eruption at the edge of the caldera. Such an eruption would melt large quantities of ice, leading to a major jokulhlaup, accompanied by ash fall.

Other scenarios cannot be excluded.

- **From the Icelandic Met Office:** The Aviation Colour Code for Bardarbunga remains at 'orange'.
- Next meeting of The Scientific Advisory Board will be held on Monday, 29. September, unless deemed necessary.

The National Commissioner of the Icelandic Police, Department of Civil Protection and Emergency Management  
[www.almannavarnir.is](http://www.almannavarnir.is) [www.avd.is](http://www.avd.is) Twitter: [@almannavarnir](https://twitter.com/almannavarnir)