



**NATIONAL COMMISSIONER OF THE ICELANDIC POLICE**  
DEPARTMENT OF CIVIL PROTECTION AND EMERGENCY MANAGEMENT



## THE SCIENTIFIC ADVISORY BOARD OF THE ICELANDIC CIVIL PROTECTION

**Date:** 08.12.2014    **Time:** 09:30    **Location:** Crisis Coordination Centre, Skogarhlid.

**Regarding:** Volcanic activity in the Bardarbunga system.

**Attending:** Scientists from Icelandic Met Office and the Institute of Earth Sciences University of Iceland along with representatives from the Icelandic Civil Protection, the Environmental Agency of Iceland and the Directorate of Health.

### Main points

- Volcanic eruption in Holuhraun
- Air quality
- Scenarios

### Notes

- Insubstantial changes have been in the volcanic eruption in Holuhraun over the last few weeks. Tomorrow, Tuesday 9. December, will be 100 days from its outbreak.
- Seismic activity in Bardarbunga continues to be strong. The strongest earthquake since noon on Friday, 5. December, was magnitude M5,1 on Friday at 21:05. About 20 earthquakes between M4,0-5,0 were detected since Friday and 10 earthquakes between magnitude M3,0-4,0. In total about 300 earthquakes were detected in Bardarbunga since noon on Friday.
- Little seismic activity was detected in the dyke and around the eruption site in Holuhraun since Friday.
- GPS measurements show no changes. The subsidence towards Bardarbunga continues with similar rate as has been.
- Telecommunications with the GPS station in Bardarbunga caldera have not been established yet. Due to bad weather forecast it is unlikely that scientists will be able to travel to Bardarbunga in the next few days. A connection will be established with the GPS station as soon as possible.

#### Air quality:

- Today (Monday) gas pollution is expected east and south of the eruption site before noon, but north and northwest of it in the afternoon. Tomorrow (Tuesday) the pollution is expected north and northeast of the volcano.
- The Icelandic Met Office provides two-day forecasts on gas dispersion from the eruptive site in Holuhraun. Most reliable are the forecast maps approved by meteorologist on duty, see [Gas forecast](#). And although still being developed further, an automatic forecast, see [Gas model](#), is also available (trial run, see [disclaimer](#)).
- Measurements of air quality can be found on the webpage [www.airquality.is](http://www.airquality.is) Data from handheld gas monitors, spread around the country, can also be found on that page
- Instructions:
  - People who feel discomfort are advised to stay indoors, close their windows, turn up the heat and turn off air conditioning. Use periods of good air quality to ventilate the house. People experiencing adverse



# NATIONAL COMMISSIONER OF THE ICELANDIC POLICE

## DEPARTMENT OF CIVIL PROTECTION AND EMERGENCY MANAGEMENT



effects should be in immediate contact with their healthcare centre. Measurements of air quality can be found on the webpage [www.airquality.is](http://www.airquality.is) The Meteorological Office issues forecast on its web-page and warnings if conditions change to the worse.

- Instructions from [The Environment Agency of Iceland](#) and [Chief Epidemiologist](#) can be found on their web-sites.
  - Check the Icelandic Met Office forecasts for sulphuric gas dispersion on the web as described above.
  - Handheld meters have been distributed around the country for SO<sub>2</sub> measurements three times a day.
  - Information and any questions on air pollution can be sent to The Environment Agency through the email [gos@ust.is](mailto:gos@ust.is). The Environment Agency is especially looking for information from people who have been in contact with high concentrations of gas; where they were, at what time it happened, how the gas cloud looked (colour and thickness of the cloud) and how they were affected by it.
- The volcanic eruption has now been going on for over three months, the lava flow is still great in Holuhraun and the rate of the subsidence of the Bardarbunga caldera is still significant. Three scenarios are considered most likely:
    - The eruption in Holuhraun continues until the subsidence of the Bardarbunga caldera stops. The eruption can still go on for many months.
    - The volcanic fissure may lengthen southwards under Dyngjajokull, resulting in a jokulhlaup and an ash-producing eruption. It is also possible that eruptive fissures could develop in another location under the glacier. If such an eruption would be prolonged it could eventually produce a lava flow.
    - Volcanic eruption in the Bardarbunga caldera. Such an eruption would melt large quantities of ice, leading to a major jokulhlaup, accompanied by ash fall.

Other scenarios cannot be excluded.

- **From the Icelandic Met Office:** The Aviation Colour Code for Bardarbunga remains at 'orange'.
- The next meeting will be held on Monday 8 of December.

The National Commissioner of the Icelandic Police, Department of Civil Protection and Emergency Management  
[Almannavarnir](http://Almannavarnir) [www.avd.is/en](http://www.avd.is/en) Twitter: [@almannavarnir](https://twitter.com/almannavarnir)