



**NATIONAL COMMISSIONER OF THE ICELANDIC POLICE**  
DEPARTMENT OF CIVIL PROTECTION AND EMERGENCY MANAGEMENT



**THE SCIENTIFIC ADVISORY BOARD OF THE ICELANDIC CIVIL PROTECTION**

**Date: 06.10.2014**    **Time: 09:30**    **Location: Crisis Coordination Centre, Skogarhlid.**

**Regarding: Volcanic activity in the Bardarbunga system.**

**Attending: Scientists from Icelandic Met Office and the Institute of Earth Sciences University of Iceland along with representatives from the Icelandic Civil Protection, The Environment Agency of Iceland, the Directorate of Health and Vatnajökull National Park.**

**Main points**

- **Volcanic eruption in Holuhraun**
- **Air quality**
- **Scenarios**

**Notes**

- The eruption continues at a similar intensity as previous days. The lava production is steady and running to the southeast from the grader into Jökulsá á Fjöllum.
- The subsidence of the Bardarbunga caldera continues with similar rate although the size of the largest earthquakes have subsided slightly.
- Seven earthquakes greater than M3.0 were recorded since noon yesterday in or around the caldera. The largest was M5.0 at 16:58 yesterday.
- Very little seismic activity is now detected in the northern part of the dyke and around the eruption site.
- GPS measurements show minor movements.
- No change was detected in water monitoring that cannot be explained by changing weather.
- Earthquakes have been recorded near Kópasker over the last few days. Six earthquakes greater the M2,0 were recorded over the last 48 hours. The largest was M3,4 at 9:34 yesterday.

**Air quality:**

- Today (Monday) and tomorrow (Tuesday) the gas pollution from the eruption is expected to be distributed towards west and southwest. A map showing the gas forecast can be found on the web page of the Icelandic Met Office [www.vedur.is/vedur/spar/textaspar/oskufok/](http://www.vedur.is/vedur/spar/textaspar/oskufok/) An interactive map showing the gas distribution can be seen at [www.vedur.is/vedur/spar/gasdreifing](http://www.vedur.is/vedur/spar/gasdreifing)
- The Icelandic Met Office has a form on its web-page for the public to report if they have detected gas pollution. A link to the page can be found on the Icelandic version of the web page under [Skrá mengun](#).
- **Instructions:**
  - People who feel discomfort are advised to stay indoors, close their windows, turn up the heat and turn off air conditioning. Use periods of good air quality to ventilate the house. People experiencing adverse effects should be in immediate contact with their healthcare centre. Measurements of air quality can be found on the webpage [www.airquality.is](http://www.airquality.is) The Meteorological Office issues forecast on its web-page and warnings if conditions change to the worse.



# NATIONAL COMMISSIONER OF THE ICELANDIC POLICE

## DEPARTMENT OF CIVIL PROTECTION AND EMERGENCY MANAGEMENT



- Instructions from [The Environment Agency of Iceland](#) and [Chief Epidemiologist](#) can be found on their web-sites.
  - The Icelandic Met Office will publish forecasts for sulphuric gases dispersion on the web and in the national radio.
  - Information and any questions on air pollution can be sent to The Environment Agency through the email [gos@ust.is](mailto:gos@ust.is). The Environment Agency is especially looking for information from people who have been in contact with high concentrations of gas; where they were, at what time it happened, how the gas cloud looked (colour and thickness of the cloud) and how they were affected by it.
- Three scenarios are considered most likely:
    - The eruption on Holuhraun declines gradually and subsidence of the Bardarbunga caldera stops.
    - Large-scale subsidence of the caldera occurs, prolonging or strengthening the eruption on Holuhraun. In this situation, it is likely that the eruptive fissure would lengthen southwards under Dyngjufokull, resulting in a jokulhlaup and an ash-producing eruption. It is also possible that eruptive fissures could develop in another location under the glacier.
    - Large-scale subsidence of the caldera occurs, causing an eruption at the edge of the caldera. Such an eruption would melt large quantities of ice, leading to a major jokulhlaup, accompanied by ash fall.

Other scenarios cannot be excluded.

- **From the Icelandic Met Office:** The Aviation Colour Code for Bardarbunga remains at 'orange'.
- From today, the Scientific Advisory Board will meet on Mondays, Wednesdays and Fridays. The next meeting will be held on Wednesday 8 October.

The National Commissioner of the Icelandic Police, Department of Civil Protection and Emergency Management  
[Almannavarnir Civil Protection and Emergency Management](#), Twitter: [@almannavarnir](#)