



**NATIONAL COMMISSIONER OF THE ICELANDIC POLICE**  
DEPARTMENT OF CIVIL PROTECTION AND EMERGENCY MANAGEMENT



**THE SCIENTIFIC ADVISORY BOARD OF THE ICELANDIC CIVIL PROTECTION**

**Date:** 12.12.2014    **Time:** 09:30    **Location:** Crisis Coordination Centre, Skogarhlid.

**Regarding:** Volcanic activity in the Bardarbunga system.

**Attending:** Scientists from Icelandic Met Office and the Institute of Earth Sciences University of Iceland along with representatives from the Icelandic Civil Protection and the Environmental Agency of Iceland.

**Main points**

- Volcanic eruption in Holuhraun
- Air quality
- Scenarios

**Notes**

- Insubstantial changes have been in the volcanic eruption in Holuhraun over the last few weeks. Indication show that the lava is now mainly flowing inside closed channels that open near the edges of the lava field.
- Seismic activity in Bardarbunga continues to be strong. The strongest earthquake since noon on Wednesday, 10. December, was magnitude M4,6 yesterday at 12:40 on the northern rim of the caldera. About 15 earthquakes larger than M4,0 were detected since Wednesday and about 20 earthquakes between M3,0-4,0. In total about 130 earthquakes were detected in Bardarbunga since noon on Wednesday.
- Little seismic activity was detected in the dyke and around the eruption site in Holuhraun since Wednesday. About 10 earthquakes were detected during that time, all smaller than M1,1.
- GPS measurements show continuing slow subsidence towards Bardarbunga.
- Telecommunications with the GPS station in Bardarbunga caldera have not been established yet. Due to bad weather forecast it is unlikely that scientists will be able to travel to Bardarbunga in the next few days. A connection will be established with the GPS station as soon as possible.

**Air quality:**

- Today (Friday) gas pollution is expected in the south, and later in the day to the east and northeast of the eruption. Tomorrow (Saturday) gas pollution might be felt in the northeast parts.
- The Icelandic Met Office provides two-day forecasts on gas dispersion from the eruptive site in Holuhraun. Most reliable are the forecast maps approved by meteorologist on duty, see [Gas forecast](#). And although still being developed further, an automatic forecast, see [Gas model](#), is also available (trial run, see [disclaimer](#)).
- Measurements of air quality can be found on the webpage [www.airquality.is](http://www.airquality.is) Data from handheld gas monitors, spread around the country, can also be found on that page

**Instructions:**

- People who feel discomfort are advised to stay indoors, close their windows, turn up the heat and turn off air conditioning. Use periods of good air quality to ventilate the house. People experiencing adverse effects should be in immediate contact with their healthcare centre. Measurements of air quality can be found on the webpage [www.airquality.is](http://www.airquality.is) The Meteorological Office issues forecast on its web-page and warnings if



# NATIONAL COMMISSIONER OF THE ICELANDIC POLICE

## DEPARTMENT OF CIVIL PROTECTION AND EMERGENCY MANAGEMENT



conditions change to the worse.

- Instructions from [The Environment Agency of Iceland](#) and [Chief Epidemiologist](#) can be found on their websites.
  - Check the Icelandic Met Office forecasts for sulphuric gas dispersion on the web as described above.
  - Handheld meters have been distributed around the country for SO<sub>2</sub> measurements three times a day.
  - Information and any questions on air pollution can be sent to The Environment Agency through the email [gos@ust.is](mailto:gos@ust.is). The Environment Agency is especially looking for information from people who have been in contact with high concentrations of gas; where they were, at what time it happened, how the gas cloud looked (colour and thickness of the cloud) and how they were affected by it.
- The volcanic eruption has now been going on for over three months, the lava flow is still great in Holuhraun and the rate of the subsidence of the Bardarbunga caldera is still significant. Three scenarios are considered most likely:
    - The eruption in Holuhraun continues until the subsidence of the Bardarbunga caldera stops. The eruption can still go on for many months.
    - The volcanic fissure may lengthen southwards under Dyngjufokull, resulting in a jokulhlaup and an ash-producing eruption. It is also possible that eruptive fissures could develop in another location under the glacier. If such an eruption would be prolonged it could eventually produce a lava flow.
    - Volcanic eruption in the Bardarbunga caldera. Such an eruption would melt large quantities of ice, leading to a major jokulhlaup, accompanied by ash fall.

Other scenarios cannot be excluded.

- **From the Icelandic Met Office:** The Aviation Colour Code for Bardarbunga remains at 'orange'.
- The next meeting will be held on Monday 15 of December.

The National Commissioner of the Icelandic Police, Department of Civil Protection and Emergency Management

[Almannavarnir](#) [www.avd.is/en](http://www.avd.is/en) Twitter: [@almannavarnir](https://twitter.com/almannavarnir)